

West Nile Virus Disease Cases* and Presumptive Viremic Blood Donors by State – United States, 2016

State	Neuroinvasive Disease Cases [†]	Non-neuroinvasive Disease Cases	Total cases	Deaths	Presumptive Viremic Blood Donors [‡]
Alabama	13	6	19	0	4
Arizona	57	21	78	6	11
Arkansas	8	1	9	1	1
California	335	107	442	19	50
Colorado	59	90	149	8	9
Connecticut	1	0	1	0	0
District of Columbia	1	0	1	0	0
Florida	6	2	8	0	3
Georgia	4	2	6	1	6
Idaho	3	6	9	0	0
Illinois	98	56	154	8	9
Indiana	15	3	18	2	5
Iowa	16	21	37	1	5
Kansas	18	19	37	4	10
Kentucky	5	3	8	1	1
Louisiana	38	21	59	2	6
Maryland	6	0	6	0	0
Massachusetts	10	7	17	0	3
Michigan	42	1	43	3	5
Minnesota	38	45	83	5	17
Mississippi	27	16	43	2	13
Missouri	9	2	11	0	0
Montana	3	3	6	1	2
Nebraska	35	60	95	1	30
Nevada	13	3	16	0	1
New Jersey	11	0	11	2	2
New Mexico	6	0	6	1	0
New York	20	2	22	1	2
North Carolina	2	0	2	0	0
North Dakota	24	61	85	2	5
Ohio	12	5	17	4	5
Oklahoma	21	14	35	0	7
Oregon	2	2	4	0	1
Pennsylvania	12	4	16	2	3
Rhode Island	2	0	2	0	0
South Carolina	6	2	8	0	3
South Dakota	35	117	152	6	16
Tennessee	3	4	7	1	1
Texas	252	118	370	18	45
Utah	7	6	13	1	0
Vermont	2	0	2	0	0
Virginia	6	2	8	0	2
Washington	8	1	9	1	3
West Virginia	0	1	1	2	0
Wisconsin	10	3	13	0	1
Wyoming	8	3	11	0	1
Totals	1,309	840	2,149	106	288

*E.g., meningitis, encephalitis, or acute flaccid paralysis

[†]Includes confirmed and probable cases

[‡]Presumptive viremic blood donors (PVDs) are people who had no symptoms at the time of donating blood through a blood collection agency, but whose blood tested positive when screened for the presence of West Nile virus. Some PVDs develop symptoms after donation; of the 288 PVDs, 45 (16%) developed clinical illness and are also included as disease cases.